

# Grapevine Production in California

*Daniel Geisseler and William R. Horwath*


## Early Production

Grape cuttings were planted at the Mission San Diego as early as 1781, possibly even before that. In the early 19<sup>th</sup> century grapes were grown in small quantities at several missions [1].

By the time of the Gold Rush, the vineyard area had reached a few hundred acres and grapes were sold to high prices in Los Angeles and San Francisco [3, 10]. As a consequence of the lucrative prices, vineyards were planted and the area reached 56,000 acres in 1880 [10]. However, due to Pierce's Disease in the 1880s, areas in the south became unsuitable to vines and the industry was forced to move to central and northern California [10].

Despite this setback, California's vineyard area continued to expand, reaching

350,000 acres by 1910 [10]. However, demand did not grow nearly as fast and the resulting overproduction led grower prices tumble [8].



**Figure 1:** Area of bearing vineyards in California since 1920 [13].

## Wine Grapes

During prohibition, the number of active wineries in California shrunk to a dozen, which survived by producing sacramental and medicinal wines. Wine production shifted to private homes and bootleggers' establishments, which greatly affected quality. The wine grape area even increased during prohibition, reaching a peak in 1927 (Figure 1) [10]. This is no surprise considering that the prices realized peaked at \$ 82 per ton in 1921 [13]. However, the increased acreage together with the low quality of the wines led to overproduction and the prices crushed. In the 1930s, the price was generally below \$ 20 per ton [10, 13]. The number of wineries grew quickly after prohibition was repealed and more than 400 wineries were listed in 1934 [9]. However, low quality due to poor production practices, inadequate aging, and less



**Figure 2:** Location of the five leading wine grape producing counties in California [12].

desirable grape varieties, was still an issue. With consumer demand growing slower than production, overproduction was once again inevitable <sup>[9]</sup>.

Between 1920 and 1973, the wine grape area fluctuated between 100,000 and 200,000 acres and was considerably smaller than the area for raisin production. With increasing consumer interest, the statewide acreage of wine grapes rose sharply in the 1970s, and more than doubled within 4 years, reaching 318,000 acres in 1977 <sup>[12]</sup>. The wine grape area now exceeded that of raisin grapes. The San Joaquin Valley and the emerging central coast wine-growing region contributed most to the expansion <sup>[6]</sup>. Most of the plantings were of varieties for fine red table wines. This changed the characteristics of the California wine industry, as most of the wine produces until this date was in the appetizer or dessert category. By 1978, over 83% of the total wine produced was classified as table wine <sup>[3]</sup>.

The boom continued and the number of wineries increased from approximately 330 in 1975 to almost 2,500 in 2006 <sup>[14]</sup>. The boom was associated with a shift in demand towards higher quality table wines. This trend was largely initiated by the now famous 1976 wine tasting in Paris, where several California wines were voted superior to French vintages <sup>[5, 14]</sup>.

With an increased demand, plantings increased again in the 1990s, with the fastest growth on the North and Central Coast <sup>[5]</sup>. Predominantly varietal grapes, such as 'Chardonnay', 'Zinfandel', 'Cabernet Sauvignon' and 'Merlot' were planted <sup>[5]</sup>. Between 1995 and 2002, the bearing acreage increased from 328,000 to 486,000 acres. Since then, the area has increased at a slower pace, reaching 506,000 acres in 2012 <sup>[12]</sup>. California now accounts for 90% of all U.S. wine production <sup>[15]</sup>.

Despite the shift of wine production to the coastal area, San Joaquin was still the County




Figure 3: Grapevine yields since 1920 in California <sup>[13]</sup>.

with the largest area of bearing wine grapes. In 2012, the 109,500 acres accounted for 22% of the area in California. Other important wine grape counties were Fresno, Sonoma, Napa and Monterey, which had a share of 9 to 12% each (Figure 2) <sup>[12]</sup>. Wine grape production differs considerably between the coastal regions and the inland regions. In the coastal regions, production of high-quality wine dominates, while in the inland regions lower-quality wines sold in jugs or boxes at low prices are prevalent <sup>[14]</sup>. This division is also reflected in the yields. The average yield in 2011 and 2012 in Fresno and San Joaquin County was 9.5 tons/acre, while it reached only 3.6 tons in Napa and Sonoma County <sup>[2]</sup>. Overall, the wine grape yield averages some 7 tons/acre (Figure 3) <sup>[12]</sup>.

In 2012, 38% of the bearing acreage was planted to white wine varieties, with 'Chardonnay' being the by far most popular variety <sup>[13]</sup>. 'Chardonnay' production has increased tremendously since the 1970s, among other reasons because it is a versatile grape that can grow in a wide range of soil and climatic conditions and is also popular with consumers <sup>[14]</sup>. The most popular red wine varieties were 'Cabernet Sauvignon', 'Zinfandel', 'Merlot', and 'Pinot Noir' <sup>[13]</sup>.

## Raisin Grapes

Since the early 1920, the area of bearing raisin grapes has remained relatively constant, fluctuating between 200 and 300 thousand acres<sup>[12]</sup>. In 2012, the acreage of bearing raisin grapes was 205,000 acres in California. 72% of the area was located in Fresno County, followed by Madera County which had a share of 14% (Figure 4)<sup>[12]</sup>.

In the 1940, the vineyards of California produced 40% of the world's raisins<sup>[10]</sup>. In 2011/12, virtually all U.S. raisins produced in California, and the U.S. raisin production accounted for 30% of the world production, with<sup>[7, 11]</sup>

The first raisin variety, 'Muscat of Alexandria' was introduced in 1861 and was first planted in the Fresno area in 1873 on 25 acres<sup>[10]</sup>. 'Thompson Seedless' was introduced in 1872 and soon became the most popular variety<sup>[10]</sup>. Today, 'Thomson Seedless' is still the dominant variety, accounting for approximately 90% of California's raisin production<sup>[13]</sup>.



**Figure 4:** Location of the two leading raisin grape producing counties in California<sup>[12]</sup>.

The market trends for raisin production are not completely decoupled from the trends in the wine and table grape market, as 'Thompson Seedless' grapes are used as table grapes and can be crushed for the production of lower-quality wine products<sup>[4, 6]</sup>.

## Table Grapes

California. The area has been relatively constant since the 1920s, fluctuating between 60,000 and 120,000 acres<sup>[12]</sup>.

Initially produced mainly for local consumption, shipments of table grapes to the Eastern U.S. increased in the early parts of the 20<sup>th</sup> century thanks to increased demand and improved transportation<sup>[10]</sup>. Today, 98% of the U.S. table grapes are produced in California<sup>[7]</sup>.

Table grape production is concentrated in the southern San Joaquin Valley, with 85% of the harvested area being located in Kern and Tulare County in 2012 (Figure 5)<sup>[12]</sup>. A number of varieties are planted, with 'Crimson Seedless' and 'Flame Seedless' currently being the most popular<sup>[13]</sup>.



**Figure 5:** Location of the two leading table grape producing counties in California<sup>[12]</sup>.

## References

1. Adams, F., 1946. Historical background. In: Hutchison, C.B. (Ed.) California Agriculture. University of California Press, Berkeley and Los Angeles. pp. 1-50.
2. Agricultural Commissioner. Annual Crop Report. Available online at: <http://www.co.fresno.ca.us/DepartmentPage.aspx?id=54231>; <http://www.co.san-joaquin.ca.us/agcomm/annualrpts.aspx>; [http://www.sonoma-county.org/agcomm/crop\\_report.htm](http://www.sonoma-county.org/agcomm/crop_report.htm); <http://www.countyofnapa.org/AgCommissioner/CropReport/>.
3. Brown, D.S., Webb, A.D., 1983. Fruit and nut crops – Grapes. In: A.F. Scheuring (Ed.) A Guidebook to California Agriculture. University of California Press, Berkeley and Los Angeles. pp. 147-152.
4. Christensen, L.P. 2000. Raisin grape varieties. In: Raisin Production Manual. University of California, Agricultural and Natural Resources Publication 3393, Oakland, CA. Pages 38-47. Available online at: [iv.ucdavis.edu/files/24430.pdf](http://iv.ucdavis.edu/files/24430.pdf)
5. Heien, D.M. Martin, P.L., 2003. California's wine industry enters new era. California Agriculture 57(3), 71-75.
6. Johnston, W.E., McCalla, A.F., 2004. Whither California agriculture: Up, down, or out? Some thoughts about the future. Special Report Series, Giannini Foundation of Agricultural Economics, UC Berkeley. Available online at: <http://escholarship.org/uc/item/4232w2sr>
7. Johnston, W.E., 2003. Cross sections of a diverse agriculture. In: Siebert, J. (Ed.) California Agriculture: Dimensions and Issues. Information Series Giannini Foundation of Agricultural Economics, UC Berkeley. pp 29-55. Available online at: <http://www.escholarship.org/uc/item/9145n8m1>
8. Moulton, K.S., 1980. Wine – a multibillion-dollar industry. California Agriculture 34(7), 9-11.
9. Singleton, V.L., Berg, H.W., Boulton, R.B., Webb, A.D., 1980. A century of wine and grape research. California Agriculture 34(7), 4-5.
10. Tufts, W.P., Briggs, F.N., Conrad, J.P., Gilmore, J.W., Hendry, G.W., Mackie, W.W., Madson, B.A., Smith, F.L., 1946. The rich pattern of California crops. In: Hutchison, C.B. (Ed.) California Agriculture. University of California Press, Berkeley and Los Angeles. pp. 113-238.
11. USDA FAS, 2012. Raisins: World markets and trade. Available online at: [usda01.library.cornell.edu/usda/fas/raiswm//2010s/2012/raiswm-09-14-2012.pdf](http://usda01.library.cornell.edu/usda/fas/raiswm//2010s/2012/raiswm-09-14-2012.pdf)
12. USDA/NASS, 2013a. Quickstats. Available online at <http://quickstats.nass.usda.gov/>
13. USDA/NASS, 2013b. Grape acreage report. [http://www.nass.usda.gov/Statistics\\_by\\_State/California/Publications/Grape\\_Acreage/](http://www.nass.usda.gov/Statistics_by_State/California/Publications/Grape_Acreage/)
14. Volpe, R.J. III, Green, R., Heien, D., Howitt, R., 2010. Wine-grape production trends reflect evolving consumer demand over 30 years. California Agriculture 64(1), 42-46.
15. The Wine Institute, 2013. California wine profile 2012. Available online at: [www.wineinstitute.org/files/CA\\_Wine\\_Stat\\_Profile\\_2012.pdf](http://www.wineinstitute.org/files/CA_Wine_Stat_Profile_2012.pdf)

---

Daniel Geisseler is a post-doctoral scientist in the Department of Land, Air and Water Resources at the University of California, Davis.

William R. Horwath is professor of Soils and Biogeochemistry in the Department of Land, Air and Water Resources and the James G. Boswell Endowed Chair in Soil Science at the University of California, Davis.

The document has been prepared within the project "Assessment of Plant Fertility and Fertilizer Requirements for Agricultural Crops in California", funded by the California Department of Food and Agriculture Fertilizer Research and Education Program (FREP).

This document is available online at [http://apps.cdfa.ca.gov/frep/docs/Grapevine\\_Production\\_CA.pdf](http://apps.cdfa.ca.gov/frep/docs/Grapevine_Production_CA.pdf)

Last update: April, 2014